

**GCE**

**Chemistry B**

**H433/02: Scientific literacy in chemistry**

Advanced GCE

**Mark Scheme for Autumn 2021**

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of candidates of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, Cambridge Nationals, Cambridge Technicals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support, which keep pace with the changing needs of today's society.




This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by examiners. It does not indicate the details of the discussions which took place at an examiners' meeting before marking commenced.

All examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the report on the examination.



© OCR 2021


## 1. Annotations


Annotation	Meaning
	Correct response
	Incorrect response
	Omission mark
<b>BOD</b>	Benefit of doubt given
<b>CON</b>	Contradiction
<b>RE</b>	Rounding error
<b>SF</b>	Error in number of significant figures
<b>ECF</b>	Error carried forward
<b>L1</b>	Level 1
<b>L2</b>	Level 2
<b>L3</b>	Level 3
<b>NBOD</b>	Benefit of doubt not given
<b>SEEN</b>	Noted but no credit given
<b>I</b>	Ignore

2. Abbreviations, annotations and conventions used in the detailed Mark Scheme (to include abbreviations and subject-specific conventions).

<b>Annotation</b>	<b>Meaning</b>
<b>DO NOT ALLOW</b>	Answers which are not worthy of credit
<b>IGNORE</b>	Statements which are irrelevant
<b>ALLOW</b>	Answers that can be accepted
<b>( )</b>	Words which are not essential to gain credit
<b>—</b>	Underlined words must be present in answer to score a mark
<b>ECF</b>	Error carried forward
<b>AW</b>	Alternative wording
<b>ORA</b>	Or reverse argument

Question			Answer	Mark	AO	Guidance
1	a	i	inert/unreactive ✓	1	1.2	
1	a	ii	high-boiling liquid ✓ porous support ✓	2	2 x 1.2	
1	b	i	mass spectroscopy/spectrometry ✓	1	2.7	<b>ALLOW</b> compare retention times with data book
1	b	ii	(compound) 1/ methylbenzene/ C <sub>6</sub> H <sub>5</sub> CH <sub>3</sub> ✓	1	1.1	<b>ALLOW</b> C <sub>7</sub> H <sub>8</sub>
1	c		(compound) 4/ octane ✓ Least/no branching/chains can get close (ORA) ✓ most Van der Waals forces (ORA) ✓	3	3 x 2.3	<b>ALLOW</b> London/id-id
1	d		C <sub>8</sub> H <sub>18</sub> → C <sub>4</sub> H <sub>8</sub> + C <sub>4</sub> H <sub>10</sub> ✓	1	2.5	
1	e		no/lack of rotation about C=C ✓ two different groups on each C ✓	2	2 x 1.2	<b>ALLOW</b> 2 diagrams to illustrate E/Z isomers
1	f		Bromine/solution decolorised/ goes (from brown/yellow/orange to) colourless ✓  ✓  ✓ (sulfuric) acid <b>and</b> (potassium) dichromate/formulae ✓ heat/reflux ✓	5	1.2 2.3 2.3 1.2 1.2	If starting colour of bromine is given it must be brown/yellow/orange <b>ALLOW</b> 'decolourised' on its own as bromine named in stem of Question <b>ALLOW</b> any unambiguous structural formula <b>IGNORE</b> brackets or 'n'  <b>ALLOW</b> 'OH' rather than 'O-H'
1	g		<b>CHECK ANSWER ON ANSWER LINE</b> <b>If answer is C<sub>3</sub>H<sub>8</sub> award 3 marks</b>  amount CO <sub>2</sub> = 11/44 = 0.25 mol <b>and</b> amount H <sub>2</sub> O = 6/18 = 0.33 mol ✓ Ratio = 3:4 <b>or</b> C:H 3:8 ✓ Hydrocarbon is C <sub>3</sub> H <sub>8</sub> ✓	3	3 x 2.6	<b>ALLOW</b> ecf Eg If nCO <sub>2</sub> = 0.25 <b>AND</b> nH <sub>2</sub> O = 0.3 then the formula for the Hydrocarbon is C <sub>5</sub> H <sub>12</sub> and scores 2 marks If no other marks scored a formula of C <sub>3</sub> H <sub>4</sub> scores 1 mark
				19		

Question			Answer	Mark	AO	Guidance
2	a	i	0.5/ ½ ✓ 2 ✓	2	2 x 2.6	ALLOW 'multiples'
2	a	ii	(½)O <sub>2</sub> ✓ oxidation state goes from 0 to -2 ✓	2	2 x 2.1	ALLOW 'oxygen' IGNORE 'O'
2	b	i	 <p>both electrodes and solutions ✓ salt bridge and voltmeter ✓ 1.0 mol dm<sup>-3</sup> and 298 K ✓</p>	3	3 x 3.3	IGNORE composition of salt bridge  IGNORE concentrations for first mark If concentrations not given, then '(aq)' or 'solution' must be there for first mark.
2	b	ii	0.32 (V) ✓	1	2.4	
2	c		(Student is incorrect) – electrons flow from Zn to Fe ✓  Reason – $E_{Zn}$ more negative than $E_{Fe}$ ✓ (Student is <b>incorrect</b> ) Half-equation 2 is not reversed/occurs in forward direction (as $E_{Fe}$ is more positive than $E_{Zn}$ ). ✓ (Student is incorrect) The sea-water acts as the salt bridge ✓	4	3.2 3.1 3.1 3.2	Electrons flow from more negative Zn to less negative Fe scores MP1 and MP2 <b>DO NOT ALLOW</b> ' $E_{Zn}$ less/lower than $E_{Fe}$ ' <b>ALLOW</b> half equation 1 is reversed
2	d		Fe(OH) <sub>2</sub> (s) + ¼ O <sub>2</sub> (g) + ½ H <sub>2</sub> O(l) → Fe(OH) <sub>3</sub> (s) correct formulae ✓ equation correctly balanced ✓ all ss correct ✓	3	3 x 2.7	ALLOW multiples  ALLOW even if equation/balancing incorrect
2	e		amount Fe <sub>2</sub> O <sub>3</sub> •2H <sub>2</sub> O = 5/195.6 (= 0.0256) mol ✓ mass water = 0.0256 x 2 x 18 = 0.92 g so students incorrect ✓	2	3.1 3.2	ALLOW ecf <b>OR</b> amount of Fe <sub>2</sub> O <sub>3</sub> = 4/159.6 = 0.025 ✓ Amount of water = 1/18 = 0.056, so ratio is 1:2.2, so student is incorrect ✓  <b>OR</b> amount of water = 1/18 = 0.056 mol ✓ Expected mass of Fe <sub>2</sub> O <sub>3</sub> •2H <sub>2</sub> O = 0.5 x 0.056 x 195.6 = 5.47 g, so students incorrect ✓ If incorrect Mr for Fe <sub>2</sub> O <sub>3</sub> •2H <sub>2</sub> O used leading to a calculation giving mass water = 1g then allow MP2 via ecf
				17		

Question			Answer	Mark	AO	Guidance
3	a		107: M+1 peak / C <sub>6</sub> <sup>13</sup> CH <sub>6</sub> O <sup>(+)</sup> ✓ 77: C <sub>6</sub> H <sub>5</sub> <sup>+</sup> ✓	2	2 x 2.2	Clear indication of C <sup>13</sup> within the parent ion '+' sign not essential for first mark but must be somewhere for second mark.
3	b		Rings(AW) (of electron density/delocalised electrons) ✓ above and below (plane of) benzene/carbon ring ✓	2	2 x 1.2	
3	c		benzene has (electrophilic) substitution <b>and</b> ethene has (electrophilic) addition ✓	1	2.1	
3	d	i	complementary colour/blue-violet is absorbed ✓ (Student incorrect because molecule) does not emit light ✓ yellow colour is what is left after absorption(AW) ✓	3	3 x 3.1	<b>ALLOW</b> yellow is the colour transmitted/reflected
3		ii	<b>CHECK ANSWER ON ANSWER LINE</b> <b>5.7 x 10<sup>-5</sup> (cm) scores 3 marks</b>  Use of $\lambda = hc/E$ ✓ $E = 6.63 \times 10^{-34} \times 3 \times 10^8 / 3.5 \times 10^{-19} = 5.7 \times 10^{-7} \text{ m}$ ✓ Convert to cm 5.7 x 10 <sup>-5</sup> (cm) ✓	3	3 x 2.6	<b>ALLOW</b> ecf <b>ALLOW</b> 2 or more sf <b>ALLOW</b> use of $E = h\gamma$ and $c = \lambda\gamma$ with subsequent rearrangement to calculate the value of $\lambda$ $\gamma = E/h = 3.5 \times 10^{-19} / 6.63 \times 10^{-34} = 5.28 \times 10^{14}$ ✓ $\lambda = c/\gamma = 3 \times 10^8 / 5.28 \times 10^{14} = 5.7 \times 10^{-7}$ ✓ convert to cm giving 5.7 x 10 <sup>-5</sup> ✓ Units not required but if given they must be correct.
3	e	i	 attack by CN <sup>-</sup> ✓ intermediate and attack by H <sup>+</sup> ✓	2	1.2	<b>IGNORE</b> partial charges and product Curly arrows must start (if projected) on lone pair on C (or negative charge if no lone pair) or double bond and end (if projected) on the appropriate atom <b>ALLOW</b> negative charge on either atom of cyanide ion.
3	e	ii	cyan(o)hydrin/hydroxynitrile ✓	1	1.1	

3	e	iii	<b>CHECK ANSWER ON ANSWER LINE</b> <b>Answer rounding to 88% scores 3 marks</b>  amount benzaldehyde = $5/106$ (= 0.0472 mol) ✓ amount mandelic acid = $6.3/152$ (= 0.0414 mol) ✓ yield = $0.0414 \times 100/0.0472 = 88\%$ ✓ OR Amount benzaldehyde = $5/107$ (=0.0467) ✓ Amount mandelic acid = $6.3/153$ (=0.0412) ✓ Yield = $0.0412 \times 100/0.0467 = 88\%$ ✓	3	3 x 2.8	<b>ALLOW</b> 88 to 2 or more sf.  Alternative method: Mr values 106 and 152 ✓ max mass of mandelic acid = $5 \times 152/106 = 7.17\text{g}$ ✓ % = $6.3 \times 100/7.17 = 88\%$ ✓ use of parent ion Mr = 107 giving Mr mandelic acid as 153 and subsequent evaluation $5 \times 153/107 = 7.15 \text{ g}$ , ✓ so % = $6.3 \times 100/7.15 = 88\%$ ✓
---	---	-----	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---	---------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------



3	f*	<p>Refer to marking instructions on page 5 of mark scheme for guidance on marking this question.</p> <p><b>Level 3 (5-6 marks)</b> Correct structure/identity for <b>both</b> A and B deduced and most evidence related to each structure is provided from <b>both</b> spectra.</p> <p><i>There is a well-developed line of reasoning which is clear and logically structured. The information presented is relevant and substantiated.</i></p> <p><b>Level 2 (3-4 marks)</b> Correctly identifies one compound with most evidence for that compound <b>OR</b> Correctly identifies both compounds with some supporting evidence</p> <p><i>There is a line of reasoning presented with some structure. The information presented is relevant and supported by some evidence.</i></p> <p><b>Level 1 (1-2 marks)</b> Correctly identifies one compound with no valid evidence <b>OR</b> Structure/identity not given or incorrect for both compounds, but some correct evidence from at least one spectrum.</p> <p><i>There is an attempt at a logical structure with a line of reasoning. The information is in the most part relevant and correct.</i></p> <p><b>Level 0 (0 marks)</b> No response or nothing worthy of credit.</p>	6	4 x 3.1 2 x 3.2	<p><b>Indicative scientific points include:</b></p> <p><b>AO3.1 Analysis:</b></p> <p><b>IR</b></p> <ul style="list-style-type: none"> <li>• <b>A</b> has OH 3200-3600 (broad)</li> <li>• <b>A</b> has no C=O around 1700</li> <li>• <b>B</b> has C=O at 1700</li> <li>• <b>B</b> has carboxylic O-H at 2500-3300 (broad)</li> <li>• Both have aromatic C=C peaks in range 1450-1650</li> </ul> <p><b><sup>13</sup>C-NMR</b></p> <ul style="list-style-type: none"> <li>• <b>A</b> has aromatic carbons 120 – 140</li> <li>• <b>A</b> has C-O at 65</li> <li>• <b>B</b> has aromatic carbons 'around 130' (AW)</li> <li>• <b>B</b> has C=O at 175</li> </ul> <p>If no relevant analysis given in text look at the spectra for any identification of bonds.</p> <p><b>AO3.2 Evaluation</b></p> <ul style="list-style-type: none"> <li>• <b>A</b> is C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>OH/phenylmethanol/correct structure drawn</li> <li>• <b>B</b> is C<sub>6</sub>H<sub>5</sub>COOH/benzoic acid/correct structure drawn</li> </ul> <p><b>IGNORE</b> references to number of C atoms(given in question)</p> <p><b>IGNORE</b> remarks relating structure of A and B to benzaldehyde (question asks for analysis of spectra)</p>
			23		

Question			Answer	Mark	AO	Guidance
4	a	i	measure water using measuring cylinder ✓ insulated cup ✓ measure initial temp ✓  add solid and measure final temp <u>when all dissolved/when temperature (change) becomes constant</u> ✓	4	3.4 3.4 3.3 3.3	Eg polystyrene, polythene, styrofoam etc. (must be clear that this is before the addition of the solid)
4	a	ii	<b>CHECK ANSWER ON ANSWER LINE +33 (kJ mol<sup>-1</sup>) scores 4 marks</b>  150 x 4.18 x 5.3 = 3.3(23) kJ/3323 J ✓ 10.1/101 = 0.1 mol ✓ 3.3(23)/0.1 = 33 kJ mol <sup>-1</sup> ✓ 2sf and '+' sign ✓	4	2.8 2.8 2.8 3.1	Allow any number of sf for MP1 to MP3  1. calculates Q using mcθ 2. calculate amount of KNO <sub>3</sub> 3. converts to kJ per mol (any sign or no sign) 4. converts to 2sf <b>and</b> includes '+' sign
4	b	i	Entropy is no. of ways of arranging particles / a measure of randomness/disorder / associated energy quanta ✓ (KIO <sub>3</sub> ) solid/ (ionic) lattice has lower entropy than (ions in) solution (ora) ✓	2	2 x 2.7	<b>ALLOW</b> +ve sign shows that particles <u>in solution</u> are in a <u>more random arrangement</u> than in a solid with a higher entropy for 2 marks  +ve sign shows that particles have become more disordered/random for 1 mark
4	b	ii	Use of $\Delta_{\text{tot}}S = \Delta_{\text{sys}}S - \Delta H/T$ ✓ Converts $\Delta H$ from kJ to J ✓ calculates $\Delta_{\text{tot}}S = 0$ at 434K ✓ Student is correct/reaction is not feasible as $\Delta_{\text{tot}}S$ would be negative below this temp/434K ✓	4	4 x 3.1	Recall equation  Insert values and evaluates correctly <b>ALLOW</b> reaction is not feasible if T < 434 since $\Delta_{\text{tot}}S$ will have a -ve value
4	c	i	same number/two of moles (of gas) on each side (of the equation) ✓	1	2.6	

4	c*	<p>ii Refer to marking instructions on page 5 of mark scheme for guidance on marking this question.</p> <p><b>Level 3 (5-6 marks)</b> Explains how temperature and pressure affect both rate and yield, giving most fine detail and makes at least one related recommendation about the process.</p> <p><i>There is a well-developed line of reasoning which is clear and logically structured. The information presented is relevant and substantiated.</i></p> <p><b>Level 2 (3-4 marks)</b> Explains how rate <b>or</b> yield is affected by both temperature <b>and</b> pressure, with some fine detail, and makes at least one recommendation about the process <b>or</b> Explains how both rate <b>and</b> yield, are affected by either temperature <b>or</b> pressure, with some fine detail, and makes one related recommendation about the process</p> <p><i>There is a line of reasoning presented with some structure. The information presented is relevant and supported by some evidence.</i></p> <p><b>Level 1 (1-2 marks)</b> Attempts to explain how rate and yield are affected by temperature or pressure <b>or</b> Attempts to explain how rate or yield is affected by both temperature and pressure</p> <p><i>There is an attempt at a logical structure with a line of reasoning. The information is in the most part relevant.</i></p> <p><b>Level 0 (0 marks)</b> No response or nothing worthy of credit.</p>	6	<p>1.2 x 2 3.1 x 1 3.2 x 3</p>	<p><b>Indicative scientific points may include:</b> <i>(fine detail in italic)</i></p> <p><b>AO 1.2 Knowledge and Understanding</b></p> <ul style="list-style-type: none"> <li>• high temperature increases rate/ORR</li> <li>• <i>As a greater frequency of collisions have <math>E \geq E_A</math></i></li> <li>• rate would be higher at higher P/ORR</li> <li>• <i>collisions are more frequent</i></li> </ul> <p><b>AO3.1 Analysis</b></p> <ul style="list-style-type: none"> <li>• A higher temperature would give a greater yield</li> <li>• <i>Reaction is endothermic</i></li> <li>• Yield does not depend on pressure</li> <li>• <i>Equal moles on each side (AW)</i></li> <li>• <i>explanations in terms of Le Chatelier</i></li> </ul> <p><b>AO3.2 Recommendations</b></p> <ul style="list-style-type: none"> <li>• Suggests optimum conditions of high temperature and low pressure (Approx. 1 – 2 atm and 500K) to maximise yield</li> <li>• <i>high temperature would be limited by cost/safety (AW)</i></li> <li>• <i>high pressure would be limited by cost/safety (AW).</i></li> </ul>
			21		

Question			Answer	Mark	AO	Guidance
5	a	i	f-(block)	1	2.1	
5	b	i	Bright/ coloured lines on a black background ✓	1	1.2	
5	b	ii	when electrons fall energy levels, energy is released ✓ $\Delta E = hv$ / energy proportional to frequency ✓ electrons in elements have different energy levels – different lines in spectrum ✓	3	1.2 1.2 2.5	
	c	i	104.5 $\pm$ 1 ✓	1	1.2	
5	c	ii	Four areas of electron density (around O) ✓  repel as far apart as possible/ minimise repulsion ✓ (two) lone pairs repel more than bonding pairs ✓	3	3x 1.2	<b>ALLOW</b> four electron pairs (around O) – can be shown on a suitable diagram OR as a description such as 2 bp and 2 lp.
5	d		Both have covalent bonding Si–O ✓  Gorilla has (in addition) Al–O bonds ✓ and ionic bonds ✓	3	3 x 3.1	<b>ALLOW</b> both based on Silicon Dioxide/contain silicon and oxygen <b>IGNORE</b> references to amorphous/crystalline <b>ALLOW</b> Gorilla contains aluminosilicate anions / sodium ions / potassium ions
5	e		Both form 1+ ions / ions have the same charge ✓ K has more electrons/shells than Na ✓	2	2 x 1.1	
5	f		Al: 2 x +3 <b>and</b> Si: 14 x +4 = 62 ✓ O: 32 x 2 = 64, so x = (-)2 ✓	2	2 x 2.2	
5	g	i	In <sup>3+</sup> (since In in Group 13/3) ✓ gives In <sub>2</sub> O <sub>3</sub> with O <sup>2-</sup> ✓	2	2 x 2.2	<b>ALLOW</b> any reasoned explanation using oxidation states or ionic charges
	g	ii	In:Si:O :: 53/114.8: 28/118.7; 19/16 :: 0.46:0.24:1.19 ✓ :: 2: 1: 5 is SnO <sub>2</sub> + In <sub>2</sub> O <sub>3</sub> ✓	2	2 x 2.6	<b>IGNORE</b> In <sub>2</sub> SnO <sub>5</sub>
				20		

**OCR (Oxford Cambridge and RSA Examinations)**  
**The Triangle Building**  
**Shaftesbury Road**  
**Cambridge**  
**CB2 8EA**

**OCR Customer Contact Centre**

**Education and Learning**

Telephone: 01223 553998

Facsimile: 01223 552627

Email: [general.qualifications@ocr.org.uk](mailto:general.qualifications@ocr.org.uk)

[www.ocr.org.uk](http://www.ocr.org.uk)

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored